

The averaging of Hamiltonian structures in discrete variant of Whitham method

A.Ya.Maltsev

L.D.Landau Institute for Theoretical Physics, e-mail: maltsev@itp.ac.ru

In this work we shall construct the procedure of averaging of Hamiltonian structures in discrete variant of Whitham method. This procedure will be analogous to Dubrovin-Novikov procedure of averaging of local field-theoretical brackets. In discrete variant of Whitham method we obtain the evolution of slow modulated m-phase solutions of nonlinear systems of type:

$$\dot{\varphi}_n^i = f^i(\varphi_n, \varphi_{n-1}, \varphi_{n+1}, \dots), \quad (1)$$

defined on the space of fields $\varphi = (\varphi_n^1, \dots, \varphi_n^p)$, where n runs all integer numbers and functions f depend on finite number of arguments.

Quasi-periodic m-phase solutions of system (1) are the solutions of the form:

$$\varphi_n^i(t) = \Phi^i(\mathbf{k}(\mathbf{U})n + \omega(\mathbf{U})t + \theta_0, \mathbf{U}), \quad (2)$$

where $\mathbf{k}(\mathbf{U})$ and $\omega(\mathbf{U})$ are m-dimensional vector, $\mathbf{U} = (U^1, \dots, U^N)$ are parameters of solution. Here 2π -periodic functions of m variables (θ) $\Phi^i(\theta, \mathbf{U})$ satisfy to the conditions:

$$\omega^\alpha(\mathbf{U})\Phi_{\theta^\alpha}^i(\theta, \mathbf{U}) = f^i(\Phi(\theta, \mathbf{U}), \Phi(\theta - \mathbf{k}(\mathbf{U}), \mathbf{U}), \Phi(\theta + \mathbf{k}(\mathbf{U}), \mathbf{U}), \dots). \quad (3)$$

Let system (1) is Hamiltonian with respect to Poisson bracket:

$$\{\varphi_n^i, \varphi_k^j\} = \sum_m B_{(m)}^{ij}(\varphi_n, \varphi_{n-1}, \varphi_{n+1}, \dots) \delta_{n,k-m}, \quad (4)$$

where the sum contains a finite number of terms (the description of such bracket can be found in [3]), with Hamiltonian function:

$$H[\varphi] = \sum_{m=-\infty}^{\infty} \mathcal{P}_H(\varphi_n, \varphi_{n-1}, \varphi_{n+1}, \dots) \quad (5)$$

and has N commutative conservation laws of the form:

$$I^\nu = \sum_{n=-\infty}^{\infty} \mathcal{P}^\nu(\varphi_n, \varphi_{n-1}, \varphi_{n+1}, \dots), \quad \nu = 1, \dots, N, \quad (6)$$

$$\{I^\nu, I^\mu\} = 0, \quad \{I^\nu, H\} = 0 \quad (7)$$

($B_{(m)}^{ij}$, $\mathcal{P}_H$ and $\mathcal{P}^\nu$ depend on finite number of arguments).

We shall have view (1):

$$\frac{d}{dt}\mathcal{P}^\nu(\varphi_n, \varphi_{n-1}, \dots) = \sum_m [Q_{(m)}^\nu(\varphi_n, \varphi_{n-1}, \dots) - Q_{(m)}^\nu(\varphi_{n-m}, \varphi_{n-m-1}, \dots)]$$

(there is a finite number of terms in the sum, $Q_{(m)}^\nu$ depend on finite number of arguments). Whitham system, giving the evolution of slow modulated parameters $\mathbf{U}$, can be written in the form:

$$\partial/\partial T \langle \mathcal{P}^\nu \rangle(\mathbf{U}(X)) = \sum_m m \partial/\partial X \langle Q_{(m)}^\nu \rangle(\mathbf{U}(X)), \quad (8)$$

here $\langle \dots \rangle$ means the averaging on the family (2), defined by the formula:

$$\langle F(\varphi_n, \varphi_{n-1}, \dots) \rangle(\mathbf{U}) = 1/(2\pi)^m \int F(\Phi(\theta, \mathbf{U}), \Phi(\theta - \mathbf{k}(\mathbf{U}), \mathbf{U}), \dots) d^m \theta.$$

View (4) we shall have:

$$\{\mathcal{P}^\nu(\varphi_n, \dots), \mathcal{P}^\mu(\varphi_k, \dots)\} = \sum_m A_{(m)}^{\nu\mu}(\varphi_n, \varphi_{n-1}, \dots) \delta_{n,k-m} \quad (9)$$

and, view (7),

$$\sum_m A_{(m)}^{\nu\mu}(\varphi_n, \varphi_{n-1}, \dots) = \sum_l (Q_{(l)}^{\nu\mu}(\varphi_n, \varphi_{n-1}, \dots) - Q_{(l)}^{\nu\mu}(\varphi_{n-l}, \varphi_{n-l-1}, \dots))$$

(all sums contain finite numbers of terms, $A_{(m)}^{\nu\mu}$ and $Q_{(l)}^{\nu\mu}$ depend on finite numbers of arguments).

Theorem 1. Let parameters $\mathbf{U}$ are such, that $U^\nu = \langle \mathcal{P}^\nu \rangle$, then:

1) System (8) is Hamiltonian with respect to Poisson bracket of Hydrodynamic Type (see [4]-[5]), defined by the relations:

$$\{U^\nu(X), U^\mu(Y)\} = \sum_m m \langle A_{(m)}^{\nu\mu} \rangle(X) \delta'(X-Y) + \left[\frac{\partial}{\partial X} \sum_l l \langle Q_{(l)}^{\nu\mu} \rangle(X) \right] \delta(X-Y) \quad (10)$$

with Hamiltonian function: $H = \int \langle \mathcal{P}_H \rangle(X) dX$.

2) From (10) follow the relations:

$$\{k^\alpha(\mathbf{U}(X)), U^\nu(Y)\} = \omega^{\alpha(\nu)}(\mathbf{U}(X)) \delta'(X-Y) + \frac{\partial \omega^{\alpha(\nu)}(\mathbf{U}(X))}{\partial X} \delta(X-Y),$$

$$\{k^\alpha(\mathbf{U}(X)), k^\beta(\mathbf{U}(Y))\} = 0, \quad \alpha, \beta = 1, \dots, m, \quad \nu = 1, \dots, N, \quad (11)$$

where $\omega^{(\nu)}(\mathbf{U})$ are frequencies analogous to ω and corresponding to flows generated by integrals I^ν on the family of solutions (2).

Theorem 2. Let system (1) has two different sets of integrals of form (6), $\{I^1, \dots, I^N\}$ and $\{\bar{I}^1, \dots, \bar{I}^N\}$, possessing all necessary properties. Then brackets (10) defined with the aid of these sets coincide with each other. That is, if on the family (2) we have: $U^\nu = I^\nu$, $\bar{U}^\nu = \bar{I}^\nu$, then bracket (10), defined with the aid of the set $\{I^1, \dots, I^N\}$, transforms into the bracket defined with the aid of the set $\{\bar{I}^1, \dots, \bar{I}^N\}$ after the transformation of parameters $U^\nu = U^\nu(\bar{\mathbf{U}})$ of m-phase solutions (2).

The proof of Theorems 1 and 2 is analogous to the proof of the same theorems for the averaged by Dubrovin-Novikov method field-theoretical brackets ([6],[7]) and is based on Dirac restriction of Poisson bracket:

$$\{\varphi^i(\theta, X), \varphi^j(\theta', Y)\} = \sum_m B_{(m)}^{ij}(\varphi(X), \varphi(X - \epsilon), \dots) \delta(X - Y + m\epsilon) \delta(\theta - \theta')$$

on the sub-manifold of functions, such that $\varphi(\theta, X)$ at any X is the function from (3) (and we use the expansion $\delta(X - Y + m\epsilon) = \delta(X - Y) + m\epsilon\delta'(X - Y) + \dots$ in the convolution with smooth functions.) The necessary conditions of regularity of this sub-manifold and the coordinates used for restriction are described in [6].

The author is grateful to S.P.Novikov and O.I.Mokhov for the attention to the work.

The work was done under the financial support of Russian Foundation for Fundamental Research (grant 96-01-01623) and INTAS (grant INTAS 96-0770).

References.

- [1] G. Whitham // Linear and Nonlinear waves. Wiley, New York, 1974.
- [2] Luke J.C. // Proc. Roy. Soc. London Ser. A, 1966. V. 292. No. 1430, 403-412.
- [3] B.A.Dubrovin // Functional Analysis and Its Appl., 1989. Vol. 23, N. 2, P. 57-59.
- [4] B.A.Dubrovin, S.P.Novikov // Dokl. Akad. Nauk SSSR, 1983. Vol. 270, N. 4, P. 781-785.
- [5] B.A.Dubrovin, S.P.Novikov // Uspekhi Mat. Nauk (Russian Mat. Surv.), 1989. Vol. 44, N. 6, P. 29-98.

- [6] A.Ya.Maltsev. //”The conservation of the Hamiltonian structures in Whitham’s method of averaging”. solv-int/9611008, 1996 (to appear) .
- [7] A.Ya.Maltsev. // Uspekhi Mat. Nauk. 1997. Vol. 52, N. 2, P. 177-178.